

Going in circles looking
for migraine relief?

Elyxyb[™]
(celecoxib)
Oral Solution



**ELYXYB: AN ORAL SOLUTION FOR A MIGRAINE
WITH OR WITHOUT AURA IN ADULTS.**

It can be painful searching for the appropriate medication.
It's time for a different oral solution.



ELYXYB
Head in a new direction.

APPROVED USE

ELYXYB[™] (celecoxib) oral solution is a prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine.
- It is not known if ELYXYB is safe and effective in children.

IMPORTANT SAFETY INFORMATION about ELYXYB[™]

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

ELYXYB™

An oral solution for migraine.

ELYXYB is the first and only ready-to-use oral liquid solution for the acute treatment of migraine with or without aura in adults.



Formulated for fast onset

ELYXYB works within 1 hour for most patients, even for moderate to severe migraine. It works to treat your migraine attack, regardless of when you take it.*



Long-lasting

Many patients in clinical trials reported sustained pain relief from 2 hours for up to 24 hours after taking ELYXYB.*



Safety in clinical studies

No drowsiness was reported and no serious adverse events occurred during clinical studies.* Please see page 7 for possible side effects with ELYXYB.

*As shown in clinical studies vs placebo.

Important safety information about ELYXYB:

ELYXYB contains celecoxib, a NSAID. NSAIDs, including ELYXYB, can cause serious side effects, including:

- **Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:**
 - o with increasing doses of NSAIDs
 - o with longer use of NSAIDs

Do not take ELYXYB right before or after a heart surgery called a “coronary artery bypass graft (CABG).”

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.



Big relief. Small bottle.



ACTUAL SIZE

A 2.5-inch bottle

That contains a single dose of medication that may help relieve migraine. No premixing, shaking, or measuring required.

A less-than-1-teaspoon dose

That can be taken on an empty stomach or with food. After your dose, you may sip up to 8 ounces of water if desired.

A take-anywhere option

That you can conveniently take on the go. No refrigeration needed.

**No prep. No steps.
Just open and drink.**

Avoid taking NSAIDs, including ELYXYB, after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- **Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines:**
 - o anytime during use
 - o without warning symptoms
 - o that may cause death

Relief within reach.



Don't plan around migraine. Reach for ELYXYB™.

For those middle-of-the-night or early morning migraine attacks

ELYXYB can be taken on an empty stomach.

For those midday migraine attacks

ELYXYB did not cause drowsiness in clinical studies.

For those who cannot swallow pills during a migraine

ELYXYB is a less-than-1-teaspoon liquid dose.

The ready-anywhere-you-need-it
migraine treatment

Important safety information (cont):

ELYXYB should only be used:

- o exactly as prescribed
- o for the shortest time needed

Do not take ELYXYB:

- if you are allergic to celecoxib or any of the ingredients in ELYXYB.
- if you are allergic to sulfonamides.
- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.



Savings made simple.

Get started with the **ELYXYB Passport savings program** today.

Commercially insured patients may pay as little as \$0 for all prescriptions with an activated copay card.*



Remove the ELYXYB Passport Copay Card attached to this page.



Activate your copay card by:

- Texting **ACTIVATE** to 96793,
- Calling 1-855-447-6042,
- Visiting **elyxyb.com/savings**, or
- Scanning the code on the back of the brochure



Present your ELYXYB Passport Copay Card to your pharmacist along with your prescription.

*For commercially insured patients only. See eligibility requirements and copay card details inside this brochure.

Patient Access Support & Savings Program

RxBIN: 610524
RxPCN: Loyalty
RxGRP: 50778048
ISSUER: (80840)
ID: XXXXXXXXXX


Elyxyb[™] (celecoxib)
Oral Solution
Passport

RxCrossroads

By McKesson

Pay as little as \$0

Please see Important Safety Information and full Prescribing Information, including Boxed Warning at elyxyb.com/PI.

APPROVED USE

ELYXYB™ (celecoxib) oral solution is a prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine.
- It is not known if ELYXYB is safe and effective in children.

IMPORTANT SAFETY INFORMATION about ELYXYB™

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

ELYXYB contains celecoxib, a NSAID. NSAIDs, including ELYXYB, can cause serious side effects, including:

- **Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:**
 - o with increasing doses of NSAIDs
 - o with longer use of NSAIDs

Do not take ELYXYB right before or after a heart surgery called a “coronary artery bypass graft (CABG).”

Avoid taking NSAIDs, including ELYXYB, after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- **Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines:**
 - o anytime during use
 - o without warning symptoms
 - o that may cause death

To the Patient: You must present this card to the pharmacist along with your prescription to participate in this program. If you have any questions regarding your eligibility or benefits, or if you wish to discontinue your participation, call the ELYXYB Patient Access Support & Savings program at 1-855-447-6042 (8:00 AM-8:00 PM EST, Monday-Friday). When you use this card, you are certifying that you understand the program rules, regulations, and terms and conditions. You are not eligible if prescriptions are paid by any state or other federally funded programs, including, but not limited to Medicare or Medicaid, or Medigap, VA or DOD or TriCare, or where prohibited by law; and you will otherwise comply with the terms above.

To the Pharmacist: When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under any federal, state or other governmental programs for his prescription.

- Submit transaction to Rx Acquisition Company d/b/a/ RxCrossroads by McKesson using BIN #610524
- If primary commercial prescription insurance exists, input card information as secondary coverage and transmit using the COB segment of the NCPDP transaction. Applicable discounts will be displayed in the transaction response.
- Acceptance of this card and your submission of claims for the ELYXYB Patient Access Support & Savings program are subject to the LoyaltyScript® program Terms and Conditions posted at www.mckesson.com/mprstnc
- Patient is not eligible if prescriptions are paid in part or full by any state or federally funded programs, including but not limited to Medicare or Medicaid, Medigap, VA, DOD or TriCare and where prohibited by law.
- The LoyaltyScript® card is not valid for use with any other prescription drug discount or cash cards for ELYXYB. Claims submitted utilizing the program are subject to audit or validation.
- **For questions regarding setup, claim transmission, patient eligibility or other issues, call the LoyaltyScript® for the ELYXYB program at 1-855-447-6042 (8:00 AM-8:00 PM EST, Monday-Friday).**

Collegium Pharmaceutical, Inc. reserves the right to rescind, revoke or amend this offer at any time.

Collegium Pharmaceutical, Inc. © 2022

ELYXYB should only be used:

- o exactly as prescribed
- o for the shortest time needed

Do not take ELYXYB:

- if you are allergic to celecoxib or any of the ingredients in ELYXYB.
- if you are allergic to sulfonamides.
- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

Don't start a new medication before talking to your healthcare provider and telling them about all of the medications you take.**Before taking ELYXYB, tell your healthcare provider about all of your medical conditions, including if you:**

- have liver or kidney problems.
- have a history of stomach ulcer or bleeding in your stomach or intestines.
- have heart disease or risk factors that increase your chance of getting heart disease.
- have high blood pressure.
- have asthma.
- are pregnant or plan to become pregnant. Taking NSAIDs, including ELYXYB, at about 20 weeks of pregnancy or later may harm your unborn baby. If you need to take NSAIDs for more than 2 days when you are between 20 and 30 weeks of pregnancy, your healthcare provider may need to monitor the amount of fluid in your womb around your baby. **You should not take NSAIDs after about 30 weeks of pregnancy.**
- are breastfeeding or plan to breastfeed. ELYXYB may pass into your breast milk. Talk with your healthcare provider about the best way to feed your baby if you take ELYXYB.

When taking ELYXYB**See the detailed “Instructions for Use” on how to take ELYXYB solution.**

- Take ELYXYB exactly as your healthcare provider tells you to take it.
- Take ELYXYB by mouth with or without food.
- Do not take more than one dose in a 24-hour period.
- Use ELYXYB for the fewest number of days a month, as needed.

The possible side effects of ELYXYB are:

liver problems including liver failure, new or worse high blood pressure, heart failure, kidney problems including kidney failure, life-threatening allergic reactions, asthma attacks in people who have asthma, life-threatening skin reactions, medication overuse headaches, low red blood cells (anemia). **Other side effects of NSAIDs include:** stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Stop taking ELYXYB and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea
- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, legs, hands and feet

Get emergency help right away if you have:

Shortness of breath or trouble breathing, chest pain, weakness in one part or side of your body, slurred speech, or swelling of the face or throat.

If you take too much ELYXYB, call your healthcare provider or get medical help right away.

These are not all the side effects associated with ELYXYB. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For more information go to dailymed.nlm.nih.gov.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

Migraine relief you want. Savings you need.



Save on your prescription with the ELYXYB™ Passport savings program:

COMMERCIALY INSURED PATIENTS MAY PAY AS LITTLE AS \$0 FOR ALL PRESCRIPTIONS with an activated copay card.*

CONTINUOUS PROGRAM SUPPORT for patients.

SCAN AND SAVE with your phone, visit elyxyb.com/savings, or text **ACTIVATE** to 96793 to activate your copay card and learn more about saving on ELYXYB.



APPROVED USE

ELYXYB™ (celecoxib) oral solution is a prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine.
- It is not known if ELYXYB is safe and effective in children.

*For commercially insured patients only. See eligibility requirements and copay card details inside this brochure.

IMPORTANT SAFETY INFORMATION about ELYXYB™

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.