

Elyxyb™
(celecoxib)
Oral Solution

Small Bottle. Big Relief.

ELYXYB is the only liquid solution
designed to deliver fast and
long-lasting migraine relief



APPROVED USE

ELYXYB™ (celecoxib) oral solution is a prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine.
- It is not known if ELYXYB is safe and effective in children.

IMPORTANT SAFETY INFORMATION about ELYXYB™

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

Works as quickly as 15 minutes.

ELYXYB is the only ready-to-use liquid solution for people who need fast relief from migraine.



Fast acting

In some patients, ELYXYB starts to work as early as 15 minutes, and many patients experience complete freedom from pain as early as 60 minutes.



Long lasting

Most patients in clinical trials reported sustained pain relief up to 24 hours after taking ELYXYB.*



Safe

No one in clinical trials stopped taking ELYXYB because of side effects.* ELYXYB was not connected to feeling drowsy or dizzy. Please see page 7 for possible side effects.

*As shown in clinical studies vs placebo.

Important safety information about ELYXYB:

ELYXYB contains celecoxib, a NSAID. NSAIDs, including ELYXYB, can cause serious side effects, including:

- **Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:**
 - o with increasing doses of NSAIDs
 - o with longer use of NSAIDs

Do not take ELYXYB right before or after a heart surgery called a “coronary artery bypass graft (CABG).”

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

No prep. No steps. Just open and drink.

The ready-anywhere-you-need-it migraine treatment.



ACTUAL SIZE

A 2.5-inch bottle

That contains a single dose of medication that may help relieve migraine. No premixing, shaking, or measuring required.

A less-than-1-teaspoon dose

That can be taken on an empty stomach or with food. After your dose, you may sip up to 8 ounces of water if desired.

A take-anywhere option

That you can conveniently take on the go. No refrigeration needed.

Avoid taking NSAIDs, including ELYXYB, after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- **Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines:**
 - o anytime during use
 - o without warning symptoms
 - o that may cause death

Elyxyb™
(celecoxib)
Oral Solution

Relief within reach.

Don't plan around migraine. Reach for ELYXYB.



If you have high blood pressure or certain heart conditions that prevent you from taking triptans

ELYXYB works differently and may be safe for you. Tell your doctor if you have heart disease, risk factors for heart disease, or if you have high blood pressure.



If you still have migraine pain while taking other treatment

If you experience breakthrough migraine pain on a different migraine product, you can take ELYXYB for fast-acting relief.



If you are dissatisfied with your current treatment

Experiencing side effects? Not getting pain relief as quickly as you need it? You're not alone. As many as 40% of people are not happy with their current migraine treatment. Don't settle. Try ELYXYB.

Important safety information about ELYXYB:

ELYXYB should only be used:

- o exactly as prescribed
- o for the shortest time needed

Do not take ELYXYB:

- if you are allergic to celecoxib or any of the ingredients in ELYXYB.
- if you are allergic to sulfonamides.
- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

Savings made simple.

Get started with the **ELYXYB Passport** savings program today.

If you are commercially insured, you will pay
\$0 for your first prescription with a copay card.*



Remove the ELYXYB Passport Copay Card attached to this page, or visit **ELYXYB.com/Savings** to download.



Present your ELYXYB Passport Copay Card to your pharmacist along with your prescription.

*For commercially insured patients only. See eligibility requirements and copay card details inside this brochure.

Patient Access Support & Savings Program

Group: EC45401001
ID: 39854504969
Bin: 600426
PCN: 54

Elyxyb[™] (celecoxib)
Oral Solution
Passport

Pay as little as \$0

Powered By: **CHANGE HEALTHCARE**

Please see Important Safety Information and full Prescribing Information, including Boxed Warning at elyxyb.com/PI.

APPROVED USE

ELYXYB™ (celecoxib) oral solution is a prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine.
- It is not known if ELYXYB is safe and effective in children.

IMPORTANT SAFETY INFORMATION about ELYXYB™

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

ELYXYB contains celecoxib, a NSAID. NSAIDs, including ELYXYB, can cause serious side effects, including:

- Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:
 - o with increasing doses of NSAIDs
 - o with longer use of NSAIDs

Do not take ELYXYB right before or after a heart surgery called a “coronary artery bypass graft (CABG).”

Avoid taking NSAIDs, including ELYXYB, after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines:
 - o anytime during use
 - o without warning symptoms
 - o that may cause death

Eligible patients will pay \$0 on their first fill and \$20 for their subsequent fills of Elyxyb (celecoxib) Oral Solution. Offer valid for up to 12 uses. A valid Prescriber ID# is required on the prescription.

Patient Instructions: In order to redeem this offer you must have a valid prescription for Elyxyb. Follow the dosage instructions given by the doctor. This offer may not be redeemed for cash. By using this offer, you are certifying that you meet the eligibility criteria and will comply with the terms and conditions described in the Restrictions section below. Patients with questions about the Elyxyb Savings offer should call (1-844-414-2225).

Pharmacist: When you apply this offer, you are certifying that you have not submitted a claim for reimbursement under any federal, state, or other governmental programs for this prescription. Participation in this program must comply with all applicable laws and regulations as a pharmacy provider. By participating in this program, you are certifying that you will comply with the terms and conditions described in the Restrictions section below.

Pharmacist instructions for a patient with an Eligible Third Party: Submit the claim to the primary Third Party Payer first, then submit the balance due to Change Healthcare as a Secondary Payer COB [coordination of benefits] with patient responsibility amount and a valid Other Coverage Code, (e.g 8). The patient is responsible for \$0 on first fill and the first \$20 on subsequent fills and reimbursement will be received from Change Healthcare. Valid Other Coverage Code required. For any questions regarding Change Healthcare online processing, please call the Help Desk at (1-800-433-4893).

Restrictions: This offer is valid in the United States. Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, Tricare or other federal or state health programs (such as medical assistance programs). Cash Discount Cards and other non-insurance plans are not valid as primary under this offer. If the patient is eligible for drug benefits under any such program, the patient cannot use this offer. By using this offer, the patient certifies that he or she will comply with any terms of his or her health insurance contract requiring notification to his or her payor of the existence and/or value of this offer. It is illegal to (or offer to) sell, purchase, or trade this offer. Program expires (12/31/2023). This offer is not transferable and is limited to one offer per person. Not valid if reproduced. Void where prohibited by law. Program managed by ConnectRx on behalf of Solix Pharmaceuticals. The parties reserve the right to rescind, revoke or amend this offer without notice at any time.

ELYXYB should only be used:

- o exactly as prescribed
- o for the shortest time needed

Do not take ELYXYB:

- if you are allergic to celecoxib or any of the ingredients in ELYXYB.
- if you are allergic to sulfonamides.
- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

Don't start a new medication before talking to your healthcare provider and telling them about all of the medications you take.**Before taking ELYXYB, tell your healthcare provider about all of your medical conditions, including if you:**

- have liver or kidney problems.
- have a history of stomach ulcer or bleeding in your stomach or intestines.
- have heart disease or risk factors that increase your chance of getting heart disease.
- have high blood pressure.
- have asthma.
- are pregnant or plan to become pregnant. Taking NSAIDs, including ELYXYB, at about 20 weeks of pregnancy or later may harm your unborn baby. If you need to take NSAIDs for more than 2 days when you are between 20 and 30 weeks of pregnancy, your healthcare provider may need to monitor the amount of fluid in your womb around your baby. **You should not take NSAIDs after about 30 weeks of pregnancy.**
- are breastfeeding or plan to breastfeed. ELYXYB may pass into your breast milk. Talk with your healthcare provider about the best way to feed your baby if you take ELYXYB.

When taking ELYXYB**See the detailed "Instructions for Use" on how to take ELYXYB solution.**

- Take ELYXYB exactly as your healthcare provider tells you to take it.
- Take ELYXYB by mouth with or without food.
- Do not take more than one dose in a 24-hour period.
- Use ELYXYB for the fewest number of days a month, as needed.

The possible side effects of ELYXYB are:

liver problems including liver failure, new or worse high blood pressure, heart failure, kidney problems including kidney failure, life-threatening allergic reactions, asthma attacks in people who have asthma, life-threatening skin reactions, medication overuse headaches, low red blood cells (anemia). **Other side effects of NSAIDs include:** stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Stop taking ELYXYB and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea
- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, legs, hands and feet

Get emergency help right away if you have:

Shortness of breath or trouble breathing, chest pain, weakness in one part or side of your body, slurred speech, or swelling of the face or throat.

If you take too much ELYXYB, call your healthcare provider or get medical help right away.

These are not all the side effects associated with ELYXYB. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For more information go to dailymed.nlm.nih.gov.

Please see full Prescribing Information, including Boxed Warning accompanying this piece, or at elyxyb.com/PI.

Migraine relief you want. Savings you need.

Save on your prescription
with the **ELYXYB Passport**
savings program:

If you are commercially insured, you
will pay \$0 for your first prescription
with a copay card.*

Scan and save
with your phone, or visit
elyxyb.com/savings
to learn more about the
savings program.



*For commercially insured patients only. See eligibility requirements and copay card details inside this brochure.

IMPORTANT SAFETY INFORMATION about ELYXYB™

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- o Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use.
- o ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery.

Gastrointestinal Bleeding, Ulceration, and Perforation

- o NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events.

Please see Important Safety Information and full Prescribing Information,
including Boxed Warning accompanying this piece, or at elyxyb.com/PI.



ELYXYB™ is a trademark of SCILEX Pharmaceuticals Inc.
All other trademarks are the property of their respective owners.
© 2023 SCILEX Holding Company. All rights reserved.
ELY-00057 04/2023

Elyxyb™
(celecoxib)
Oral Solution